

Product Information

NOVACID®

7-5-36+7CaO+0.1B

NOVACID is a specialty line of fully water-soluble, acidifying NPK fertilizers. NOVACID guarantees a balanced nutrient supply, which delivers better yield and quality. The various analyses are ideal for drip-irrigated vegetables, fruit trees, and arable crops.

Drip irrigation enhances water and nutrient use efficiency and significantly reduces nitrous oxide (N₂O) emissions compared to furrow and sprinkler irrigation systems.

Product advantages

- NOVACID's acidity neutralizes and dissolves bicarbonates, preventing clogging of pipes and drippers
- Extends the lifespan of irrigation systems
- Uniform and efficient distribution of irrigation water and fertilizers
- Increased nutrient availability in the soil solution, improving nutrient uptake by the plants
- Enriched with potassium, calcium, and boron to solve multiple deficiencies with one formula
- Low in nitrogen, allowing full and independent control over the nitrogen in the fertigation program
- Promotes environmental sustainability

Specific usage

NOVACID 7-5-36+7CaO+0.1B is designed especially for irrigation water sources deficient in calcium and boron, and is adapted for the production and ripening phases to improve fruit development.

Application rates

Crop	Dosage, kg/ha/week
Soft fruits	20-40
Vegetables	50-100
Leafy vegetables	19-40
Orchards	20-40

Contact ICL or your professional advisor for specific advice or recommendations.



Product characteristics

Maximum solubility: 68 kg/100 L water (25°C)
EC value: 1.56 mS/cm (at 1 g/L)
pH value: 2.7 (at 10 g/L)
HCO₃ reduction: 21.6 mg/L
Packaging: 25 kg bag (PE)
Item code: 6107053601

Guaranteed analysis

Total nitrogen (N)	7%
Nitric nitrogen (N-NO ₃)	3.4%
Urea nitrogen (NH ₂ -N)	3.6%
Phosphorus pentoxide (P ₂ O ₅)	5%
Potassium oxide (K ₂ O)	36%
Calcium oxide (CaO)	7%
Chloride (Cl)	26%
Boron (B)	0.1%



Product Information

NOVACID®

7-5-36+7CaO+0.1B

The right product for the right phase (partial list)

Formulation for	Formulation	Contains			No Chloride	No or low Urea
		MgO	CaO	TE		
Starter phase	6-60-10				•	•
	13-40-13+TE			•		
	10-50-10+TE			•	•	
	19-19-19+3CaO		•			
General growth phase	7-23-24+6.5CaO		•			•
	12-4-34+4MgO+TE	•		•		
	12-4-36+3CaO+TE		•	•		
	21-8-21+3CaO+TE		•	•		
	10-6-21+8MgO+TE	•		•	•	•
	19-19-19+3CaO		•			
	7-14-35+3.5MgO	•		•	•	
	10-17-22+9CaO		•		•	
	12-6-28+ME			•	•	
	15-5-29+3 CaO					
Fruit setting and developing phase	16-8-25+4MgO+ME	•		•		
	10-6-21+8MgO+TE	•		•	•	•
	19-19-19+3CaO		•			
	10-17-22+9CaO		•		•	
	14-7-14+14CaO+TE		•	•	•	•
Production and ripening phase	16-8-25+4MgO+ME	•		•		
	3-10-37+4MgO+TE	•		•	•	•
	5-5-50					
	6-6-46					
	7-5-36+7CaO+0.1B		•	•		•
	12-4-34+4MgO+TE	•		•		
	12-4-36+3CaO+TE		•	•		
	10-10-40+TE		•		•	•
	10-6-21+8MgO+TE	•		•	•	•
	12-7-32 + 4.5 CaO + TE		•	•		
	7-14-35+3.5MgO+TE	•		•	•	
	12-6-28+ME			•	•	
	15-5-29+3 CaO		•			

